Injury Frequency Associated with Motorcycle Crashes



Richard Alcorta, MD, FACEP

Maryland Institute for
Emergency Medical Services Systems



Overview

- Severity Scoring
- Injury Types by Body Area
- Matching Mechanism to injury
- Mandatory Implementation of Single Protective Device: Helmet

^{*} Maryland Trauma Registry Calendar year 2005



Injury Severity Score (ISS)


Baker 1974

- Injury Severity Score makes possible a valid numerical description of the overall severity of injury in persons who have sustained injury to more than one area of he body
- Designed for In- Hospital determination of Probability of Mortality
- Based on assessment of known anatomical injuries
- Autopsy specific data.



Maryland Trauma Registry Data

Out of the 1064 MC
 victims entering
 Maryland's Trauma
 System, Majority are in the "low acuity" 1-12 ISS



Injury with in ISS range



Length of Stay Surrogate for Severity

| Length of Stay (days) | <1 | 1 | 2-3 | 4-7 | 8- 14 | | 22- 28 | >28 | total | Avera ge LOS |
|-----------------------------|----|-----|-----|-----|----------|----|-----------|-----|-------|--------------|
| # | 21 | 485 | 214 | 197 | 84 | 30 | 12 | 21 | 1064 | 4.4 |



Matching Mechanism to Injury

- Abrasions/ Contusions
- Head and Brain Injury
- Flexion/ Extension and Axial Loading: Spine Injury
- Straddle: Pelvic Fractures



Protective Devices of Those Entering Trauma Centers

| Protective Devices | Number | Percentage |
|------------------------------|--------|------------|
| None | 205 | 19 |
| Protective Helmet | 745 | 70 |
| Padding/ Protective Clothing | 4 | 0.4 |
| Other Protective Device | 2 | 0.2 |
| Not Valued | 108 | 10 |
| Total | 1064 | |



Injury

Patients
Out of 1064 Percent

| Head Injuries **All types | 428 | 40.2 |
|---------------------------|-----|------|
| Rib Fractures | 165 | 15.5 |
| Tibia/Fibula Fractures | 154 | 14.5 |
| Thoracic Organ Injuries | 126 | 11.8 |

These are not mutually exclusive – Patient could have more than one injury type



Occipital
Fracture with
Contrecoup
Cerebral
Contusions
Hemorrhage






| Hemo/Pneumothorax | 123 | 11.6 |
|--------------------------|-----|------|
| Abdominal organ injuries | 118 | 11.1 |
| Facial fractures | 107 | 10.1 |
| Pelvis and hip fractures | 106 | 10.0 |
| Forearm fractures | 105 | 9.9 |
| Femur fractures | 99 | 9.3 |

These are not mutually exclusive –
Patient could have more than one injury type



Straddle Injury Pattern:

Pelvis fractures "open book"







| **Head Injuries with | | |
|--------------------------|----|-----|
| Hemorrhages | 87 | 8.2 |
| **Skull Fractures | 85 | 8.0 |
| Hand and Wrist Fractures | 74 | 7.0 |
| Lumbar Spine Injuries | 61 | 5.7 |
| Thoracic Spine Injuries | 52 | 4.9 |
| Humerus Fractures | 52 | 4.9 |

These are not mutually exclusive –

Patient could have more than one injury type







| Cervical Spine Injuries | 50 | 4.7 |
|----------------------------|----|-----|
| Shoulder Dislocations | 32 | 3.0 |
| Knee Dislocations | 15 | 1.4 |
| Flail Chest | 8 | 8.0 |
| Elbow Dislocations | 7 | 0.7 |
| Fractures to Sternum | 7 | 0.7 |
| Injuries to Thoracic Aorta | 5 | 0.5 |

These are not mutually exclusive – Patient could have more than one injury type





Helmet Usage and Fatality Reduction

Helmet Use Increased

■ Pre-law 24.6%

Post-law 80.5%

Number of fatalities decreased by 36.9% (1996)



Major Brain Injury R A Cowley Shock Trauma

| | MC crash Admits | Helmet Used | Helmet Not Used | Unk | Major Brain Helmet Used | Major Brain No Helmet |
|------|--------------------|----------------|-----------------------|-----|----------------------------------|--------------------------------|
| 1992 | 216 | 56 | 118 | 42 | 2% | 19% |
| 2004 | 327 | 242 | 57 | 27 | 14.9% | 26.3% |

1992 year Mandatory Helmet law went into effect.

In 2001 4.4% died with Helmets, 12.5% died without Helmets.

Helmet worn by a fatally injured motorcyclist



Inside label: Novelty Helmet not to be used on any Motorized vehicle Highway or Offroad Use

